

The following article has been accepted by the Journal of Mathematical Physics. After it is published, it will be found at <http://jmp.aip.org> .

One real function instead of the Dirac spinor function

Andrey Akhmeteli*

LTASolid Inc.

10616 Meadowglen Ln 2708

Houston, TX 77042, USA

(Dated: July 24, 2011)

Abstract

Three out of four complex components of the Dirac spinor can be algebraically eliminated from the Dirac equation (if some linear combination of electromagnetic fields does not vanish), yielding a partial differential equation of the fourth order for the remaining complex component. This equation is generally equivalent to the Dirac equation. Furthermore, following Schrödinger (Nature, **169**, 538 (1952)), the remaining component can be made real by a gauge transform, thus extending to the Dirac field the Schrödinger's conclusion that charged fields do not necessarily require complex representation. One of the two resulting real equations for the real function describes current conservation and can be obtained from the Maxwell equations in spinor electrodynamics (the Dirac-Maxwell electrodynamics). As the Dirac equation is one of the most fundamental, these results both belong in textbooks and can be used for development of new efficient methods and algorithms of quantum chemistry.

PACS numbers: 03.65.Pm;03.65.Ta;12.20.-m;03.50.De

*Electronic address: akhmeteli@ltasolid.com; URL: <http://www.akhmeteli.org>

I. INTRODUCTION

The Dirac equation is a most fundamental equation, crucial for high energy physics and significant even for such mundane devices as lead-acid batteries (Ref. [1]). It is well-known that two components of the Dirac spinor can be algebraically eliminated from the Dirac equation (Ref. [2], p.445). However, to the best of this author's knowledge, it was not noticed before that another component can be algebraically eliminated from the resulting equations, yielding a partial differential equation of the fourth order for the remaining component. While it was noted in Ref. [3] (p.28) that "the set of four first-order equations comprising the Dirac equation is generally equivalent to, and may be reduced to, a single fourth order equation", according to the authors of Ref. [3], this remark relates to currently known rigorous solutions of the Dirac equation, and this result had not been established for the Dirac equation with arbitrary given electromagnetic field (V.G. Bagrov, private communication).

It is possible to make one step further and rewrite the Dirac equation in terms of just one real function, rather than a complex function, using Schrödinger's approach, who noted (Ref. [4]) that for each solution of the equations of scalar electrodynamics (the Klein-Gordon-Maxwell electrodynamics) there is a physically equivalent (i.e. coinciding with it up to a gauge transform) solution with a real matter field, despite the widespread belief about charged fields requiring complex representation. Schrödinger concludes his work (Ref. [4]) with the following: "One is interested in what happens when [the Klein-Gordon equation] is replaced by Dirac's wave equation of 1927, or other first-order equations. This and the bearing on Dirac's 1951 theory will be discussed more fully elsewhere." To the best of this author's knowledge, Schrödinger did not publish any sequel to Ref. [4], and the lack of extension to the Dirac equation may explain the fact that Schrödinger's work did not get the attention it deserves. Such an extension is proposed here, as the only complex component of the fourth order partial differential equation equivalent to the Dirac equation can be made real in the same way. Thus, the Dirac equation can be rewritten as an equation for one real function, rather than for a Dirac spinor. This result is both important on its own and can be used for development of new efficient methods and algorithms of quantum chemistry.

II. ELIMINATION OF COMPONENTS FROM THE DIRAC EQUATION

To illustrate the parallels with beautiful but little-known Schrödinger's work [4], the Dirac equation is considered as part of spinor electrodynamics, although most derivations are valid without any changes for the Dirac equation in electromagnetic field independently of the Maxwell equations.

Schrödinger considered the equations of scalar electrodynamics:

$$(\partial^\mu + ieA^\mu)(\partial_\mu + ieA_\mu)\psi + m^2\psi = 0, \quad (1)$$

$$\square A_\mu - A_{,\nu\mu}^\nu = j_\mu, \quad (2)$$

$$j_\mu = ie(\psi^*\psi_{,\mu} - \psi_{,\mu}^*\psi) - 2e^2 A_\mu\psi^*\psi. \quad (3)$$

He observed that the complex charged matter field ψ can be made real by a gauge transform (at least locally), although it is generally believed that complex functions are required to describe charged fields.

The equations of motion in the relevant gauge (unitary gauge (Refs. [5],[6])) for the transformed 4-potential of electromagnetic field B^μ and real matter field φ are as follows:

$$\square\varphi - (e^2 B^\mu B_\mu - m^2)\varphi = 0, \quad (4)$$

$$\square B_\mu - B_{,\nu\mu}^\nu = j_\mu, \quad (5)$$

$$j_\mu = -2e^2 B_\mu\varphi^2. \quad (6)$$

Work [4] has another unique feature. While the initial Klein-Gordon equation (1) actually contains two equations for the real and imaginary components of the complex field ψ , the relevant equation (4) for the real field φ contains just one equation. What happened to the other equation? The missing equation is equivalent to the current conservation equation, and the latter can be derived from the Maxwell equations (5), as the divergence of the antisymmetric tensor vanishes.

It turns out, however, that Schrödinger's results also hold in the case of spinor electrodynamics. In general, four complex components of the Dirac spinor function cannot be made real by a single gauge transform, but three complex components out of four can be eliminated from the Dirac equation in a general case, yielding a fourth-order partial differential equation for the remaining component, which component can be made real by a

gauge transform. The resulting two equations for one real component can be replaced by one equation plus the current conservation equation, and the latter can be derived from the Maxwell equations.

Spinor electrodynamics is a more realistic theory than scalar electrodynamics, so it seems important that the charged field of spinor electrodynamics can also be described by one real function. It is not clear if similar results can be obtained for the Standard Model.

The resulting system of equations for the electromagnetic field and the real matter field is overdetermined. This fact suggests that it may be feasible to eliminate the matter field altogether, as this was done in the case of scalar electrodynamics (Ref. [7]).

Let us start with the equations of (non-second-quantized) spinor electrodynamics:

$$(i\not{\partial} - \not{A})\psi = \psi, \quad (7)$$

$$\square A_\mu - A_{,\nu\mu}^\nu = e^2 \bar{\psi} \gamma_\mu \psi, \quad (8)$$

where, e.g., $\not{A} = A_\mu \gamma^\mu$ (the Feynman slash notation). For the sake of simplicity, a system of units is used where $\hbar = c = m = 1$, and the electric charge e is included in A_μ ($eA_\mu \rightarrow A_\mu$). In the chiral representation of γ -matrices (Ref. [6])

$$\gamma^0 = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix}, \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad (9)$$

where index i runs from 1 to 3, and σ^i are the Pauli matrices. If ψ has components

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}, \quad (10)$$

the Dirac equation (7) can be written in components as follows:

$$\begin{aligned} & (A^0 + A^3)\psi_3 + (A^1 - \imath A^2)\psi_4 + \\ & + \imath(\psi_{3,3} - \imath\psi_{4,2} + \psi_{4,1} - \psi_{3,0}) = \psi_1, \end{aligned} \quad (11)$$

$$\begin{aligned} & (A^1 + \imath A^2)\psi_3 + (A^0 - A^3)\psi_4 - \\ & - \imath(\psi_{4,3} - \imath\psi_{3,2} - \psi_{3,1} + \psi_{4,0}) = \psi_2, \end{aligned} \quad (12)$$

$$\begin{aligned}
& (A^0 - A^3)\psi_1 - (A^1 - \imath A^2)\psi_2 - \\
& -\imath(\psi_{1,3} - \imath\psi_{2,2} + \psi_{2,1} + \psi_{1,0}) = \psi_3,
\end{aligned} \tag{13}$$

$$\begin{aligned}
& -(A^1 + \imath A^2)\psi_1 + (A^0 + A^3)\psi_2 + \\
& +\imath\psi_{2,3} + \psi_{1,2} - \imath(\psi_{1,1} + \psi_{2,0}) = \psi_4.
\end{aligned} \tag{14}$$

Obviously, equations (13,14) can be used to express components ψ_3, ψ_4 via ψ_1, ψ_2 and eliminate them from equations (11,12) (cf. Ref. [2], p.445). The resulting equations for ψ_1 and ψ_2 are as follows:

$$\begin{aligned}
& -\psi_{1,\mu}^{\imath\mu} + \psi_2(-\imath A_{,3}^1 - A_{,3}^2 + A_{,2}^0 + A_{,2}^3 + \imath(A_{,1}^0 + A_{,1}^3 + A_{,0}^1) + A_{,0}^2) + \\
& +\psi_1(-1 + A^\mu A_\mu - \imath A_{,\mu}^\mu + \imath A_{,3}^0 - A_{,2}^1 + A_{,1}^2 + \imath A_{,0}^3) - 2\imath A^\mu \psi_{1,\mu} = 0,
\end{aligned} \tag{15}$$

$$\begin{aligned}
& -\psi_{2,\mu}^{\imath\mu} + \imath\psi_1(A_{,3}^1 + \imath A_{,3}^2 + \imath A_{,2}^0 - \imath A_{,2}^3 + A_{,1}^0 - A_{,1}^3 + A_{,0}^1 + \imath A_{,0}^2) + \\
& +\psi_2(-1 + A^\mu A_\mu - \imath(A_{,\mu}^\mu + A_{,3}^0 + \imath A_{,2}^1 - \imath A_{,1}^2 + A_{,0}^3)) - 2\imath A^\mu \psi_{2,\mu} = 0.
\end{aligned} \tag{16}$$

As equation (15) contains ψ_2 , but not its derivatives, it can be used to express ψ_2 via ψ_1 :

$$\psi_2 = -(\imath F^1 + F^2)^{-1} (\square' + \imath F^3) \psi_1, \tag{17}$$

where $F^i = E^i + \imath H^i$, electric field E^i and magnetic field H^i are defined by the standard formulae

$$F^{\mu\nu} = A^{\nu,\mu} - A^{\mu,\nu} = \begin{pmatrix} 0 & -E^1 & -E^2 & -E^3 \\ E^1 & 0 & -H^3 & H^2 \\ E^2 & H^3 & 0 & -H^1 \\ E^3 & -H^2 & H^1 & 0 \end{pmatrix}, \tag{18}$$

and the modified d'Alembertian \square' is defined as follows:

$$\square' = \partial^\mu \partial_\mu + 2\imath A^\mu \partial_\mu + \imath A_{,\mu}^\mu - A^\mu A_\mu + 1. \tag{19}$$

Using the above notation, equation (16) can be rewritten as follows:

$$-(\square' - \imath F^3) \psi_2 - (\imath F^1 - F^2) \psi_1 = 0, \tag{20}$$

so equation(17) can be used to eliminate ψ_2 from equation (20), yielding an equation of the fourth order for ψ_1 :

$$\left((\square' - \imath F^3) (\imath F^1 + F^2)^{-1} (\square' + \imath F^3) - \imath F^1 + F^2 \right) \psi_1 = 0. \quad (21)$$

This equation is equivalent to the Dirac equation (if $\imath F^1 + F^2 \neq 0$).

It should be noted that the coefficient at ψ_2 in equation (15) is gauge-invariant (it can be expressed via electromagnetic fields). While this elimination could not be performed for zero electromagnetic fields, this does not look like a serious limitation, as in reality there always exist electromagnetic fields in the presence of charged fields, although they may be very small. However, it should be noted that free spinor field presents a special case and is not considered in this work, as it does not satisfy the equations of spinor electrodynamics. It is not clear how free field being a special case is related to the divergencies in quantum electrodynamics. It should also be noted that the above procedure could be applied to any component of the spinor function, not just to ψ_1 , yielding equations similar to equation (21). Presenting the above results in a more symmetric form is beyond the scope of this work.

While the above elimination of the third component of the Dirac spinor is straightforward, this author failed to find it elsewhere, but cannot be sure that this important result, which belongs in textbooks, was not published previously.

Using a gauge transform, it is possible to make ψ_1 real (at least locally). Then the real and the imaginary parts of equation (16) after substitution of the expression for ψ_2 will present two equations for ψ_1 . However, it is possible to construct just one equation for ψ_1 in such a way that the system containing this equation and the current conservation equation will be equivalent to equation (16).

Let us consider the current conservation equation:

$$(\bar{\psi} \gamma^\mu \psi)_{,\mu} = 0, \quad (22)$$

or

$$(\bar{\psi}_{,\mu} \gamma^\mu \psi) + (\bar{\psi} \gamma^\mu \psi_{,\mu}) = 0, \quad (23)$$

On the other hand,

$$\begin{aligned} (\bar{\psi}_{,\mu} \gamma^\mu \psi)^* &= (\bar{\psi}_{,\mu} \gamma^\mu \psi)^\dagger = \\ &= \psi^\dagger (\gamma^\mu)^\dagger (\psi_{,\mu}^\dagger \gamma^0)^\dagger = \bar{\psi} \gamma^0 \gamma^0 \gamma^\mu \gamma^0 \gamma^0 \psi_{,\mu} = \bar{\psi} \gamma^\mu \psi_{,\mu}, \end{aligned} \quad (24)$$

as $(\gamma^\mu)^\dagger = \gamma^0 \gamma^\mu \gamma^0$ (Ref. [6]), so the current conservation equation can be written as follows:

$$\begin{aligned} 2\text{Re}(\bar{\psi}\gamma^\mu\psi_{,\mu}) &= 2\text{Im}(\bar{\psi}i\cancel{\partial}\psi) = \\ &= 2\text{Im}(\bar{\psi}(i\cancel{\partial} - \cancel{A} - 1)\psi) = 0, \end{aligned} \tag{25}$$

as it is not difficult to check that values $\bar{\psi}\cancel{A}\psi$ and $\bar{\psi}\psi$ are real.

If equations (13,14,15) hold, only the second component of spinor $(i\cancel{\partial} - \cancel{A} - 1)\psi$ can be nonzero, so equation (25) can be written as follows:

$$2\text{Im}(-\psi_4^*\delta) = 0, \tag{26}$$

where δ is the left-hand side of equation (16). Therefore, if ψ_4 does not vanish identically, the system containing the current conservation equation equation (26) and the following equation

$$2\text{Re}(\psi_4^*\delta) = 0, \tag{27}$$

is equivalent to equation (16).

III. CONCLUSION

Thus, three complex components of the Dirac equation out of four can be algebraically eliminated, and the remaining component can be made real using a gauge transform. In particular, most results of work [4] for scalar electrodynamics are extended to a more realistic theory – spinor electrodynamics: in a certain gauge, the charged matter field can be described by one real function. This function satisfies a real partial differential equation (of the fourth order) and the current conservation equation, which can be obtained from the Maxwell equations. Thus, the system of equations for the function is overdetermined. As it was shown in work [7] that the matter field can be naturally eliminated from the equations of scalar electrodynamics, the results of this work suggest that the spinor field can be eliminated from spinor electrodynamics as well (and therefore spinor electrodynamics can be embedded into a quantum field theory in the same way as scalar electrodynamics (cf. Ref. [7]), but this has not been proven yet.

Acknowledgments

The author is grateful to V.G. Bagrov for his interest in this work and valuable remarks and to J. Noldus for useful discussions.

- [1] R. Ahuja, A. Blomqvist, P. Larsson, P. Pyykkö, P. Zaleski-Ejgierd , Phys. Rev. Lett. **106**, 018301(1-4) (2011).
- [2] Fayyazuddin and Riazuddin, *Quantum Mechanics* (World Scientific, Singapore, 1990).
- [3] V. G. Bagrov and D. M. Gitman, *Exact solutions of relativistic wave equations* (Kluwer, Dordrecht, 1990).
- [4] E. Schrödinger, Nature **169**, 538 (1952).
- [5] E. Deumens, Physica **18D**, 371 (1986).
- [6] C. Itzykson and J.-B. Zuber, *Quantum field theory* (McGraw-Hill, New York, 1980).
- [7] A. Akhmeteli, Int'l Journal of Quantum Information **9**, 17 (2011).